MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2019

July–September 2019—Volume 134, pp. 439–442

https://doi.org/10.5248/134.439

Pleurotheciopsis triseptata sp. nov. from China

Cong-Cong Ai[1], Ji-Wen Xia[1], Xiu-Guo Zhang[1], Li-Guo Ma[2*]

[1] *Shandong Provincial Key Laboratory for Biology of Vegetable Diseases and Insect Pests, College of Plant Protection, Shandong Agricultural University, Taian, Shandong, 271018, China*

[2] *Shandong Key Laboratory of Plant Virology, Institute of Plant Protection, Shandong Academy of Agricultural Sciences, Jinan, Shandong, 250100, China*

[*] Correspondence to: *maliguo809@163.com*

Abstract—A new anamorphic species, *Pleurotheciopsis triseptata*, is described and illustrated from specimens collected on dead stems in China. This fungus is characterized by polyblastic and sympodial conidiogenous cells producing catenate, fusiform, 3-euseptate, versicolored conidia.

Key words—microfungi, hyphomycete, taxonomy

Introduction

Pleurotheciopsis B. Sutton was erected by Sutton (1973) for *P. pusilla* (the type species) and *P. bramleyi*. The genus is characterized by distinct and brown conidiophores with polyblastic, integrated, sympodial, and denticulate conidiogenous cells that produce euseptate and hyaline to brown conidia that form acropetal unbranched chains (Sutton 1973, Ellis 1976, Castañeda & al. 2001). Assigning species in *Pleurotheciopsis* is primarily based on conidial morphological characters, including shape, septation, size, and pigmentation. Five additional species have been described (Castañeda 1985, Cazau & al. 1993, Castañeda & Iturriaga 1999, Castañeda & al. 2001, Rambelli & al. 2008), but *P. setiformis* (with synchronous and ampulliform conidiogenous cells and setae around the conidiophores) was subsequently transferred to *Ampullicephala* by R.F. Castañeda & al. (2009). Castañeda & al. (2001) provided a key to five of the accepted *Pleurotheciopsis* species.

During a survey on wood-inhabiting microfungi in China, a new species that belongs to the genus *Pleurotheciopsis* was found. Specimens are deposited in the Herbarium of Department of Plant Pathology, Shandong Agricultural University, Taian, Shandong, China (HSAUP) and the Mycological Herbarium, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China (HMAS).

Taxonomy

Pleurotheciopsis triseptata L.G. Ma & X.G. Zhang, **sp. nov.** Fig. 1

MycoBank MB 831794

Differs from *Pleurotheciopsis sylvestris* by its 3-euseptate and smaller conidia, and from *P. bramleyi* and *P. websteri* by its 3-euseptate and versicolored conidia that are bigger than *P. bramleyi* and shorter than *P. websteri*.

Type: China, Sichuan Province, Longchi National Forest Park, on dead stems of an unidentified broadleaf tree, 22 Apr. 2012, L.G. Ma (**Holotype**, HSAUP H2239; **isotype**, HMAS 146166).

Etymology: *triseptata*, referring to the number of conidial septa.

Colonies on the natural substratum effuse, brown to dark brown, hairy. Mycelium partly superficial, partly immersed, composed of branched, septate, pale brown, smooth hyphae, 2.0–3.5 µm diam. Conidiophores macronematous, mononematous, single, unbranched, erect, straight to flexuous, cylindrical, smooth, thick-walled, brown to dark brown, pale brown towards the apex, 4–9-septate, 105–240 µm long, 6.5–13.5 µm diam at the base, 4–6.5 µm wide at the tapered apex. Conidiogenous cells integrated, terminal, polyblastic, sympodial, brown, cylindrical, smooth, thick-walled, with 1–4 short, unthickened denticles, 29.5–60 × 4.5–6.5 µm. Conidial secession schizolytic. Conidia in distinct, unbranched, acropetal chains, fusiform, sometimes curved, thick-walled, middle cells brown and end cells pale brown to subhyaline, tapered towards the ends, 3-euseptate, intercalary conidia 24–33.5 × 5.5–7.5 µm, truncate at the ends, apical conidia 22.5–33 × 6–7.5 µm, obtuse at the apex and truncate at the base.

Comments—*Pleurotheciopsis triseptata* is most similar to *P. sylvestris* R.F. Castañeda & Iturr. in conidial shape and pigmentation, but *P. sylvestris* has conidia that are 4–7-septate and larger (29–49 × 7–8.5 µm; Castañeda & Iturriaga 1999). *Pleurotheciopsis bramleyi* B. Sutton and *P. websteri* Cazau & al. also resemble our new species, but *P. bramleyi* has conidia that are hyaline, 2–3-septate, and smaller (16.5–25 × 4.5–6.5 µm; Sutton 1973),



Fig. 1. *Pleurotheciopsis triseptata* (holotype, HSAUP H2239).
A, B. Conidiophores, conidiogenous cells, and conidia;
C. Conidiophore with conidiogenous cell; D. Conidia.

while conidia in *P. websteri* are 5-septate, hyaline, longer, and narrower (32–38 × 5–6 μm; Cazau & al. 1993).

Acknowledgments

The authors express gratitude to Dr. Jian Ma and Dr. Ze-Fen Yu for serving as pre-submission reviewers and for their valuable comments and suggestions. This project was supported by Young Talents Training Program of Shandong Academy of Agricultural Sciences (CXGC2018E04), National Key R&D Program of China (2017YFD0201700), National Natural Science Foundation of China (31400019, 31230001, 31093440), and the Ministry of Science and Technology of the People's Republic of China (2006FY120100).

Literature cited

Castañeda RF. 1985. Deuteromycotina de Cuba. Hyphomycetes III. Instituto de Investigaciones Fundamentales en Agricultura Tropical "Alejandro de Humboldt", Havana, Cuba.

Castañeda RF, Iturriaga T. 1999. A new species of *Pleurotheciopsis* from a rainforest in Venezuela. Mycotaxon 70: 63–68.

Castañeda RF, Calduch M, Garcia D, Izquierdo Z. 2001. A new species of *Pleurotheciopsis* from leaf litter. Mycotaxon 77: 1–5.

Castañeda RF, Gusmão LFP, Moraes Júnior VO de, Leão-Ferreira SM, Saikawa M, Minter DW, Stadler M. 2009. Two setose anamorphic fungi: *Ampullicephala* gen. nov. and *Venustosynnema grandiae* sp. nov. Mycotaxon 109: 275–288. https://doi.org/10.5248/109.275

Cazau MC, Arambarri AM, Cabello MN. 1993. New hyphomycetes from Santiago River. VI. (Buenos Aires Province, Argentina). Mycotaxon 46: 235–240.

Ellis MB. 1976. More dematiaceous hyphomycetes. Commonwealth Mycological Institute, Kew, Surrey, England.

Rambelli A, Venturella G, Ciccarone C. 2008. Dematiaceous hyphomycetes from Pantelleria mediterranean maquis litter. Flora Mediterranea 18: 441–467.

Sutton BC. 1973. Some hyphomycetes with holoblastic sympodial conidiogenous cells. Transactions of British Mycological Society 61(3): 417–429. https://doi.org/10.1016/S0007-1536(73)80112-3